From a Neutrino Factory to Carlsbad

Kirk T. McDonald

Princeton U.

mcdonald@puphep.princeton.edu

Workshop on the Next Generation U.S. Underground Science Facility

Carlsbad, NM, June 13, 2000

http://puhep1.princeton.edu/~mcdonald/nufact/

The Opportunity for a Neutrino Factory

- Recent evidence for neutrino mass and oscillation among neutrino types is literally a gift from the heavens.

- Systematic exploration of the physics of massive neutrino is possible using accelerator sources of neutrinos and large underground detectors separated by long baselines.

- The WIPP underground science facility is well sized and well located to host a large detector for neutrinos from a neutrino factory.

  The neutrino detector must be able to distinguish the sign of the muon from the reaction $\nu + N \to \mu + X$.

Oscillations of Massive Neutrinos

Neutrinos could have a small mass (Pauli, Fermi, Majorana, 1930's).

Massive neutrinos can mix (Pontecorvo, 1957).

In the example of only two massive neutrinos (that don't decay), with mass eigenstates ν_1 and ν_2 with mass difference Δm and mixing angle θ, the flavor eigenstates ν_a and ν_b are related by

$$\begin{pmatrix} \nu_a \\ \nu_b \end{pmatrix} = \begin{pmatrix} \cos\theta & \sin\theta \\ -\sin\theta & \cos\theta \end{pmatrix} \begin{pmatrix} \nu_1 \\ \nu_2 \end{pmatrix}.$$

The probability that a neutrino of flavor ν_a and energy E appears as flavor ν_b after traversing distance L in vacuum is

$$P(\nu_a \to \nu_b) = \sin^2 2\theta \sin^2 \left(\frac{1.27 \Delta m^2 [\mathrm{eV}^2]\ L[\mathrm{km}]}{E[\mathrm{GeV}]} \right).$$

The probability that ν_a does not disappear is

$$P(\nu_a \to \nu_a) = \cos^2 2\theta \sin^2 \left(\frac{1.27 \Delta m^2 [\mathrm{eV}^2]\ L[\mathrm{km}]}{E[\mathrm{GeV}]} \right).$$

A Sketch of Current Data

- The **Atmospheric Neutrino "Anomaly"** suggests that GeV ν_μ's (from $p+N_2 \to \pi \to \mu\nu_\mu$) disappear while traversing the Earth's diameter, $\Rightarrow \Delta m^2 \approx 10^{-3}$ (eV)^2 for $\sin^2 2\theta \approx 1$. (Kamiokande, IMB, Soudan-2, MACRO, Super-Kamiokande)









- The **Solar Neutrino "Deficit"** suggests that MeV ν_e's disappear between the center of the Sun and the Earth.
  $\Rightarrow \Delta m^2 \approx 10^{-10}$ (eV)^2 for $\sin^2 2\theta \approx 1$, if vacuum oscillations.
  (Homestake, Super-Kamiokande, GALLEX, SAGE)









- The **LSND Experiment** suggests that 30-MeV $\overline{\nu}_\mu$'s (from $p + H_2O \to \pi^- \to \mu^- \overline{\nu}_\mu$) appear as $\overline{\nu}_e$'s after 30 m. $\Rightarrow \Delta m^2 \approx 1\ (\mathrm{eV})^2$, but reactor data requires $\sin^2 2\theta \lesssim 0.03$.







- The atmospheric neutrino anomaly + the solar neutrino deficit (if both correct) require at least 3 massive neutrinos.

- If LSND is correct as well, need at least 4 massive neutrinos.

- The measured width of the Z^0 boson (LEP) $\Rightarrow$ only 3 Standard Model neutrinos. A 4th massive neutrino must be "sterile".

Mixing of Three Neutrinos

$$\begin{pmatrix} \nu_e \\ \nu_\mu \\ \nu_\tau \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & c_{23} & s_{23} \\ 0 & -s_{23} & c_{23} \end{pmatrix} \begin{pmatrix} c_{13} & 0 & s_{13}e^{i\delta} \\ 0 & 1 & 0 \\ -s_{13}e^{-i\delta} & 0 & c_{13} \end{pmatrix} \begin{pmatrix} c_{12} & s_{12} & 0 \\ -s_{12} & c_{12} & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} \nu_1 \\ \nu_2 \\ \nu_3 \end{pmatrix},$$

where $c_{12} = \cos\theta_{12}$, *etc.* (Maki, Nakagawa, Sakata, 1962).

Three massive neutrinos $\Rightarrow$ six independent parameters:

- Three mixing angles: θ_{12}, θ_{13}, θ_{23},
- A phase δ related to CP violation,
- Two differences of the squares of the neutrino masses.
  Ex: $\Delta m^2_{12} = \Delta m^2(\text{solar})$ and $\Delta m^2_{23} = \Delta m^2(\text{atmospheric})$.
- $[J_{CP} = s_{12}s_{23}s_{31}c_{12}c_{23}c^2_{31}s_\delta =$ Jarlskog invariant.]

Measurement of these parameters is a primary goal of experimental neutrino physics.

If four massive neutrinos, then 6 mixing angles, 3 phases, 3 independent squares of mass differences.

Matter Effects

ν_e's can interact with electrons via both W and Z^0 exchanges, but other neutrinos can only interact with e's via Z^0 exchange.

$$\Rightarrow \sin^2 2\theta_{\mathrm{matter}} = \frac{\sin^2 2\theta_{\mathrm{vac}}}{\sin^2 2\theta_{\mathrm{vac}} + (\cos 2\theta_{\mathrm{vac}} - A)^2},$$

where $A = 2\sqrt{2} G_F N_e E / \Delta m^2$ depends on sign of Δm^2.

At the "resonance", $\cos 2\theta_{\mathrm{vac}} = A$, $\sin^2 2\theta_{\mathrm{matter}} = 1$ even if $\sin^2 2\theta_{\mathrm{vac}}$ is small (Wolfenstein, 1978, Mikheyev, Smirnov, 1986).

$\Rightarrow$ 3 MSW solutions to the solar neutrino problem:



In all of these MSW solutions, $\Delta m^2_{\mathrm{solar}} = \Delta m^2_{12} > 0$.

Too Many Solutions

There are 8 scenarios suggested by present data:

- Either 3 or 4 massive neutrinos.
- Four solutions to the solar neutrino problem:
  1. Vacuum oscillation (VO, or "Just So") solution;
     $\Delta m^2_{12} \approx (0.5 - 5.0) \times 10^{-10}$ eV^2, $\sin^2 2\theta_{12} \approx (0.7 - 1.0)$.
  2. Low MSW solution;
     $\Delta m^2_{12} \approx (0.5 - 2.0) \times 10^{-7}$ eV^2, $\sin^2 2\theta_{12} \approx (0.9 - 1.0)$.
  3. Small mixing angle (SMA) MSW solution;
     $\Delta m^2_{12} \approx (4.0 - 9.0) \times 10^{-6}$ eV^2, $\sin^2 2\theta_{12} \approx (0.001 - 0.01)$.
  4. Large mixing angle (LMA) MSW solution;
     $\Delta m^2_{12} \approx (0.2 - 2.0) \times 10^{-4}$ eV^2, $\sin^2 \theta_{12} \approx (0.65 - 0.96)$.
- Atmospheric neutrino data $\Rightarrow$ $\Delta m^2_{23} \approx (3 - 5) \times 10^{-4}$ eV^2, $\sin^2 \theta_{12} > 0.8$.
- θ_{13} very poorly known ($\sin^2 2\theta_{13} \lesssim 0.2$).
- δ completely unknown.

The Next Generation of Neutrino Experiments

- Short baseline accelerator experiments (miniBoone, ORLAND, CERN) will likely clarify the LSND result.

- Super-Kamiokande + new long baseline accelerator experiments (K2K, Minos, NGS) will firm up measurements of θ_{23} and Δm^2_{23}, but will provide little information on θ_{13} and δ.

- New solar neutrino experiments (BOREXino, SNO, HELLAZ, HERON, ...) will explore different portions of the energy spectrum, and clarify possible pathlength-dependent effects.

  SNO should provide independent confirmation of neutrino oscillations via comparison of reactions
  $\nu + {}^2\text{H} \to p + p + e$ and $\nu + {}^2\text{H} \to p + n + \nu$.

- Each of these experiments studies oscillations of only a single pair of neutrinos.

- The continued search for the neutrinoless double-beta decay ${}^{78}\text{Ge} \to {}^{78}\text{Se} + 2e^-$ will improve the mass limits on Majorana neutrinos to perhaps as low as 0.01 eV (GENIUS).

The Opportunity for a Neutrino Factory

- Many of the neutrino oscillation solutions permit study of the couplings between 2, 3, and 4 neutrinos in accelerator based experiments.

- More neutrinos are needed!

- Present neutrino beams come from $\pi, K \to \mu\nu_\mu$ with small admixtures of $\overline{\nu}_\mu$ and ν_e from μ and $K \to 3\pi$ decays.





- Cleaner spectra and comparable fluxes of ν_e and ν_μ desirable.

A Neutrino Factory based on a Muon Storage Ring

- Higher (per proton beam power) and better characterized, neutrino fluxes are obtained from μ decay.

- Collect low-energy μ's from π decay,
  Cool the muon bunch,
  Accelerate the μ's to the desired energy,
  Store them in a ring while they decay via $\mu^- \to e^- \nu_\mu \overline{\nu}_e$.
  [Of course, can use μ^+ also.]




A Neutrino Factory
based on a
Muon Storage Ring
p Source,
8-24 GeV,
1-4MW
20-T Capture Solenoid
Target
Mini Cooling (LH_2)
First Phase
Rotation (rf)
Drift
2nd Phase Rotation
(Induction Linac)
2-GeV Superconducting Linac
Ionization Cooling
(LH_2, rf, Alternating
Solenoids)
2 - 50 GeV Recirculating
Linac
50-GeV Muon Storage Ring
~ 1km circumference
Neutrino Beam to Near Detector
Neutrino Beam to Far Detector
99-12-10 • Peter Gruber, CERN-PS
p
π
π+μ
μ
μ+ν
ν

6 Classes of Experiments at a Neutrino Factory

$$\nu_\mu \to \nu_e \to e^- \qquad \text{(appearance)}, \tag{1}$$

$$\nu_\mu \to \nu_\mu \to \mu^- \qquad \text{(disappearance)}, \tag{2}$$

$$\nu_\mu \to \nu_\tau \to \tau^- \qquad \text{(appearance)}, \tag{3}$$

$$\overline{\nu}_e \to \overline{\nu}_e \to e^+ \qquad \text{(disappearance)}, \tag{4}$$

$$\overline{\nu}_e \to \overline{\nu}_\mu \to \mu^+ \qquad \text{(appearance)}, \tag{5}$$

$$\overline{\nu}_e \to \overline{\nu}_\tau \to \tau^+ \qquad \text{(appearance)}. \tag{6}$$

[Plus 6 corresponding processes for $\overline{\nu}_\mu$ from μ^+ decay.]

Processes (2) and (5) are easiest to detect, via the final state μ.

Process (5) is noteworthy for having a "wrong-sign" μ.

Processes (3) and (6) with a final state τ require μ's of 10's of GeV.

Processes (1) and (4) with a final state electron are difficult to distinguish.

Magnetic detectors of 10's of kilotons will be required, with fine segmentation if τ's are to be measured.

Scaling Laws for Rates at a Neutrino Factory

$\sigma_\nu \propto E$; $\quad I_\nu \propto 1/(\theta)^2 \propto (E/L)^2$: $\quad$ Rate $\propto I_\nu \sigma_\nu \propto E^3/L^2$.

$\Rightarrow$ Rate $\propto E^3$ at fixed L, $\quad$ Rate $\propto 1/L^2$ at fixed E.



Neutrino oscillation probability varies with L/E,

$\Rightarrow$ Rate $\propto E$ for fixed L/E.



τ appearance suppressed at low energy. Larger $E \Rightarrow$ larger L.

The Rates are High at a Neutrino Factory

Charged current event rates per kton-yr.

($L = 732$ km)	ν_μ	$\overline{\nu}_e$
Neutrino Factory	(2×10^{20} ν_μ/yr)	
10 GeV	2200	1300
20 GeV	18,000	11,000
50 GeV	2.9×10^5	1.8×10^5
250 GeV	3.6×10^7	2.3×10^7
MINOS (WBB)		
Low energy	460	1.3
Medium energy	1440	0.9
High energy	3200	0.9

Even a low-energy neutrino factory has high rates of electron neutrino interactions.

A neutrino factory with $E_\mu \gtrsim 20$ GeV is competitive for muon neutrino interactions.

$\nu_\mu \to \nu_\mu \to \mu^-$ Disappearance



$E_\mu = 30$ GeV,
2×10^{20} μ decays,
$L = 7000$ km,
$\sin^2 2\theta_{23} = 1$.
(hep-ph/9906487)

Δm^2_{23} (eV^2)	Events (per 10 kt-yr)
0.002	2800
0.003	1200
0.004	900
0.005	1700
No Osc.	6200

$\nu_\mu \to \nu_\tau \to \tau^-$ Appearance

Δm^2_{23} (eV^2)	Events (per 10 kton-yr)
0.002	1200
0.003	1900
0.004	2000
0.005	1800

For conditions as above.

Measuring θ_{13}

Many ways:

$$P(\overline{\nu}_e \to \overline{\nu}_\mu) = \sin^2 2\theta_{13} \sin^2 \theta_{23} \sin^2 \frac{1.27\Delta m^2_{23} L}{E_\nu},$$

$$P(\overline{\nu}_e \to \overline{\nu}_\tau) = \sin^2 2\theta_{13} \cos^2 \theta_{23} \sin^2 \frac{1.27\Delta m^2_{23} L}{E_\nu},$$

$$P(\nu_\mu \to \nu_\tau) = \cos^4 \theta_{13} \sin^2 2\theta_{23} \sin^2 \frac{1.27\Delta m^2_{23} L}{E_\nu}.$$



10 kton detector,
$E_\mu = 20$ GeV,
2×10^{20} μ decays,
$L = 732$ km,
$\sin^2 2\theta_{23} = 1$,
Left: $\overline{\nu}_e \to \overline{\nu}_\mu \to \mu^+$,
Right: $\nu_\mu \to \nu_\mu \to \mu^-$,
Box = presently allowed.
(hep-ph/9811390).

Measuring the Sign of Δm^2_{23} via Matter Effects



The matter-effect resonance depends on the sign of Δm^2_{23} (p. 8).

Crisper to observe $\overline{\nu}_e \to \overline{\nu}_\mu$ appearance than ν_μ disappearance.





Baseline of 5000-7000 km needed to build up the matter effect for measurement of Δm^2_{23}.

[Measurement of CP violation more favorable at 3000 km.]

Measuring δ via CP Violation

The phase δ is accessible to terrestrial experiment in the large mixing angle (LMA) solution to the solar neutrino problem (or if there are 4 massive neutrinos).

CP violation:

$$A_{\rm CP} = \frac{P(\nu_e \to \nu_\mu) - P(\overline{\nu}_e \to \overline{\nu}_\mu)}{P(\nu_e \to \nu_\mu) + P(\overline{\nu}_e \to \overline{\nu}_\mu)} \approx \left| \frac{2\sin\delta}{\sin 2\theta_{13}} \sin \frac{1.27 \Delta m^2_{12} L}{E} \right|,$$

assuming $\sin^2 2\theta_{12} \approx \sin^2 2\theta_{23} \approx 1$ (LMA).



10 kton detector,
2×10^{21} muon decays,
Large angle MSW:
$\Delta m^2_{12} = 10^{-4}$ eV^2,
$\Delta m^2_{23} = 2.8 \times 10^{-3}$ eV^2,
$\theta_{12} = 22.5°$,
$\theta_{13} = 13°$,
$\theta_{23} = 45°$,
$\delta = -90°$.
(hep-ph/9909254)

Matter effects dominate the asymmetry for $L > 1000$ km.

Measuring δ via T Violation

If the small mixing angle (SMA) solutions holds, may still be able to measure δ via T violation:

$$P(\nu_e \to \nu_\mu) - P(\nu_\mu \to \nu_e) =$$

$$4J_{\rm CP} \sin \frac{1.27\Delta m^2_{12} L}{E} \sin \frac{1.27\Delta m^2_{13} L}{E} \sin \frac{1.27\Delta m^2_{23} L}{E},$$

$$8J_{\rm CP} = \cos\theta_{13} \sin 2\theta_{13} \sin 2\theta_{12} \sin 2\theta_{23} \sin\delta = \text{Jarlskog invariant}.$$

Matter effects could make $\sin 2\theta_{12}$ resonate for $E \approx 100$ MeV and $L \approx 10,000$ km (hep-ph/9911258).







However, not easy to measure $\nu_\mu \to \nu_e \to e^-$ (appearance) against background of $\overline{\nu}_e \to \overline{\nu}_e \to e^+$ in a large, massive detector in which the electrons shower immediately. [Rates low also.]

A Neutrino Factory is a Global Facility



- Host lab with the muon storage ring and near detector.
- Could have two larger detectors located elsewhere, possibly one on the same, and the second on another continent.
- For this, the muon storage ring needs 3 straight sections, and would not lie in a horizontal plane.



Large Underground Detectors

K2K:



"Spaghetti" detector concept:



Gran Sasso in Italy:

DOE WIPP facility in New Mexico:



A Neutrino Factory is a Step to a Muon Collider



A First Muon Collider to study light-Higgs production:



Summary

- The physics program of a neutrino factory/muon collider is extremely diverse, and of scope to justify an international laboratory.
- The first step is a **neutrino factory** capable of systematic exploration of neutrino oscillations.
  - With $\gtrsim 10^{20}$ ν's/year can go well beyond other existing or planned accelerator experiments.
  - Beams with $E_{\nu_e} \lesssim 1$ GeV are already very interesting.
  - Higher energy is favored: Rate $\propto E$ at fixed L/E; ν_τ appearance practical only for $E \gtrsim 30$ GeV.
  - Detectors at multiple distances needed for broad coverage of parameter space $\Rightarrow$ triangle or "bowtie" storage rings.
  - CP and T violation accessible with $\gtrsim 10^{21}$ ν's/year.
  - Control of muon polarization extremely useful when studying $\nu_e \to e$ modes.


AND WHEN THE ENTIRE UNIVERSE COMES TO A HALT, AND ALL THAT'S LEFT IS AN INCREASINGLY SCATTERED ARRAY OF ELECTRONS, POSITRONS, AND NEUTRINOS, WILL YOU LOVE ME?
I THINK WE SHOULD BE ALLOWED TO SEE OTHER NEUTRINOS.