# A Simple 3-Body Gravitational Problem

Lee H. McDonald
*3461 East 3rd Street, Tucson, AZ 85716*
Kirk T. McDonald
*Joseph Henry Laboratories, Princeton University, Princeton, NJ 08544*
(August 6, 2023)

## 1 Problem

In general, there is no analytic solution for the motion of a 3-body gravitational problem. Here is a simple version that permits an analytic solution.

Three equal masses $m$ are initially at rest along the $x$-axis with separations $R$. At time $t = 0$ the "outer" two masses become free to move, towards the center mass that remains at rest at $x = 0$.

## 2 Solution

For $t > 0$ the equation of motion of each of the "outer" accelerate according to Newton's law of gravitation with acceleration of magnitude $a$ related by,

$$ma = \frac{Gm^2}{r^2} + \frac{Gm^2}{4r^2}\,, \qquad a = \frac{dv}{dt} = \frac{5Gm}{4r^2}\,, \tag{1}$$

where $G$ is Newton's gravitational constant, $r$ is the distance between each of the two "outer" masses and the "center" mass, and $v$ is the magnitude of the velocity of each of the "outer" masses.

A first integral of eq. (1) is obtained from conservation of energy $U$,

$$\begin{aligned} U &= \mathrm{KE} + \mathrm{PE} = U_0 = -2\frac{Gm^2}{R} - \frac{Gm^2}{2R} = -\frac{5Gm^2}{2R} \\ &= U(r) = 2\frac{mv^2}{2} - 2\frac{Gm^2}{r} - \frac{Gm^2}{2r} = mv^2 - \frac{5Gm^2}{2r}\,. \end{aligned} \tag{2}$$

Hence,

$$v = \frac{dr}{dt} = \sqrt{\frac{5Gm}{2}}\sqrt{\frac{1}{r} - \frac{1}{R}}, \qquad dr\sqrt{\frac{r}{R-r}} = \sqrt{\frac{5Gm}{2R}}dt \tag{3}$$

With $x = R - r$, $x$ goes from $R$ to 0 as time increases from 0 to $t$ when the masses collide. Using 195.01 and 195.04 of [1] with $a = 0$, $b = 1$, $f = R$, $g = -1$ and $k = ag - bf = -R$,

$$\begin{aligned} \sqrt{\frac{5Gm}{2R}}t &= -\int_R^0 dx\,\sqrt{\frac{R-x}{x}} = \int_0^R dx\,\sqrt{\frac{R-x}{x}} = \sqrt{x}\sqrt{R-x}\Big|_0^R + \frac{R}{2}\int_0^R \frac{dx}{\sqrt{x}\sqrt{R-x}} \\ &= -\frac{R}{2}\sin^{-1}\frac{R-x}{R}\Big|_0^R = \frac{\pi R}{4}\,, \end{aligned} \tag{4}$$

so the collision occurs at time,

$$t = \frac{\pi R}{4}\sqrt{\frac{2R}{5Gm}}\,. \tag{5}$$

# A Appendix: Another Simple Example

The present example is closely related to the first considerations of the gravitational 3-body problem by Euler in 1767 [2, 3]. Another simple example is the case that two masses $m$ are somehow held fixed at $x = \pm R$ along the $x$-axis, and mass $M$ is near the origin. The equilibrium point for mass $M$ is the origin, and this equilibrium is stable against small perturbations.

The gravitational potential in the $x$-$y$ planes for mass $M$ is,

$$\begin{aligned} U(x,y) &= -\frac{GMm}{\sqrt{(R+x)^2+y^2}} - \frac{GMm}{\sqrt{(R+x)^2+y^2}} \\ &\approx U(0,0) + \frac{1}{2}\frac{\partial^2 U(0,0)}{\partial x^2}x^2 + \frac{1}{2}\frac{\partial^2 U(0,0)}{\partial y^2}y^2 + \frac{\partial^2 U(0,0)}{\partial x \partial y}xy. \end{aligned} \tag{6}$$

The terms in the potential in $x^2$ and $y^2$ imply that the angular frequencies for small oscillations of $M$ about the origin are,

$$\omega_x = \sqrt{\frac{\partial^2 U(0,0)/\partial x^2}{M}} = \sqrt{\frac{8Gm}{R^3}}\,, \qquad \omega_y = \sqrt{\frac{\partial^2 U(0,0)/\partial y^2}{M}} = \sqrt{\frac{2Gm}{R^3}} = \frac{\omega_x}{2}\,. \tag{7}$$

For small oscillations in $z$, $\omega_z = \omega_y$.

Note that if mass $M$ were in a circular orbit of radius $R$ about fixed mass $m$, its angular velocity would be $\Omega = \sqrt{Gm/R^3}$ (Kepler's 3$^{\rm rd}$ law).

The motion for small oscillations of the form $x = \cos(\omega_x t)$ and $y = \sin(\omega_y t)$ is sketched below.



## References


[1] H.B. Dwight, *Tables of Integrals and Other Mathematical Data*, 4$^{\rm th}$ ed. (Macmillan, 1961), http://kirkmcd.princeton.edu/examples/EM/dwight_57.pdf

[2] L. Euler, *De motu rectilineo trium corporum se mutuo attrahentium*, Novi Comm. Acad. Sci. Petrop. **11**, 144 (1767), E327,
http://kirkmcd.princeton.edu/examples/mechanics/euler_ncasp_11_144_67.pdf
https://scholarlycommons.pacific.edu/euler-works/327/

[3] S.R. Bistafa, *Annotated Translations of Three of the Euler's Papers on Celestial Mechanics*, Adv. Hist. Stud. **8**, 252 (2019),
http://kirkmcd.princeton.edu/examples/mechanics/bistafa_ahs_8_252_19.pdf